



STATUS OF FERMILAB G-2 ANALYSIS TOWARDS FINAL RESULT



SIMON CORRODI
Argonne National Laboratory

on behalf of the Muon g-2 collaboration
III Workshop on Muon Precision
Physics (MPP2024)
November 13th 2024



INTRINSIC MAGNETIC MOMENT

Magnetic moment $\vec{\mu}$ is connected to spin \vec{s} via dimensionless factor g

$$\vec{\mu} = g \frac{q}{2m} \vec{s}$$


"gyromagnetic ratio"

INTRINSIC MAGNETIC MOMENT

Magnetic moment $\vec{\mu}$ is connected to spin \vec{s} via dimensionless factor g

$$\vec{\mu} = g \frac{q}{2m} \vec{s}$$

"gyromagnetic ratio"



THE MAGNETIC MOMENT OF THE MUON: HISTORY

Storage Ring


Dilated lifetime measurement of a_μ , more precise

Stopped Muons

Stop muons in a magnetic field measurement of g_μ directly



Experiment



THE MAGNETIC MOMENT OF THE MUON: HISTORY

Storage Ring


Dilated lifetime measurement of a_μ , more precise

Stopped Muons

Stop muons in a magnetic field measurement of g_μ directly

Experiment

$\sim 3.5\sigma$ sigma between a_μ^{exp} and a_μ^{SM}



THE MAGNETIC MOMENT OF THE MUON: HISTORY


FNAL goal: 4 x improvement

Storage Ring

Dilated lifetime
measurement of a_μ , more precise

Stopped Muons


Stop muons in a magnetic field
measurement of g_μ directly



FERMILAB MUON G-2 DATA TAKING


6 years of data taking, passed the TDR goal 21 x BNL statistics

Post-Run-6 with magnet on but no muons (magnetic field syst.)




FERMILAB MUON C


6 years of data taking



Run-1 Result: 2021 :: 0.46 ppm




$$a_\mu(\text{FNAL}) = 116\,592\,040(54) \times 10^{-11} \quad (0.46 \text{ ppm})$$




FERMILAB MUON G-2 DATA TAKING

6 years of data taking, passed the TDR goal 21 x BNL statistics




FERMILAB MUON G-2 DATA TAKING

6 years of data taking, passed the TD



Run-2/3 Result: 2023 :: 0.2 ppm




$$a_\mu(\text{Exp}) = 116\,592\,059(22) \times 10^{-11} \text{ [190 ppb]}$$

Analyzed positrons (Billions)

FERMILAB MUON G-2 DATA TAKING


6 years of data taking, passed the TDR goal 21 x BNL statistics






Muon Campus

MC-1






Video by M. Fertl and R. Reimann,
Diorama: Fermi National Accelerator Laboratory



- Accelerating protons to 8 GeV
- Form 16 120 ns-long bunches
- Pion production in fixed target
- Pion decay to muons (95% polarization)
- $p = 3.094 \text{ GeV}/c \pm 5 \%$
- $\sim 10000 \mu^+$ per bunch
- Muons outrun protons


Video by M. Fertl and R. Reimann,
Diorama: Fermi National Accelerator Laboratory





- Accelerating protons to 8 GeV
- Form 16 120 ns-long bunches
- Pion production in fixed target
- Pion decay to muons (95% polarization)
- $p = 3.094 \text{ GeV}/c \pm 5 \%$
- $\sim 10000 \mu^+$ per bunch
- Muons outrun protons
- Muon g-2 experimental hall

Video by M. Fertl and R. Reimann,
Diorama: Fermi National Accelerator Laboratory

Muon g-2 at Fermilab: MC-1









polarized muons






store polarized muons
in a **dipole B** field



store polarized muons
in a **dipole B** field




store polarized muons
in a **dipole B** field

$$\omega_C = \frac{e}{m\gamma}B$$

$$\omega_S = \frac{e}{m\gamma}B(1 + \gamma a_\mu)$$

store polarized muons
in a **dipole B** field




$$\omega_C = \frac{e}{m\gamma} B$$

$$\omega_S = \frac{e}{m\gamma} B(1 + \gamma a_\mu)$$

store polarized muons
in a **dipole B field**

$$g = 2$$
$$a_\mu = 0$$

—→ momentum
—→ spin



$$\omega_C = \frac{e}{m\gamma} B$$

$$\omega_S = \frac{e}{m\gamma} B(1 + \gamma a_\mu)$$


$$\omega_a = \omega_S - \omega_C$$

store polarized muons
in a **dipole B field**

$g = 2$
 $a_\mu = 0$

$g > 2$
 $a_\mu > 0$

—→ momentum
—→ spin





The image shows a large, circular particle detector, likely a muon g-2 experiment, with a complex arrangement of equipment and a central area containing several tall, dark cabinets. A dashed blue line traces the inner boundary of the detector's structure. A red arrow points from the equation to the innermost part of the detector, and a blue arrow points from the text to the outer ring. Two workers are visible on the floor of the detector, providing a sense of scale.

$$\omega_a = \omega_s - \omega_c$$

store polarized muons
in a **dipole B** field


$$\omega_a = \omega_s - \omega_c$$

store polarized muons
in a **dipole B** field

extract the **muon magnetic anomaly**

$$\omega_a = a_\mu \frac{eB}{mc}$$

by measuring




WHAT IS THE MAGIC MOMENTUM?

extract the muon magnetic anomaly

$$\vec{\omega}_a = -\frac{q}{m} \vec{B} + a_\mu \vec{B}$$

by measuring


The diagram shows the equation $\vec{\omega}_a = -\frac{q}{m} \vec{B} + a_\mu \vec{B}$. The term $a_\mu \vec{B}$ is highlighted in red. An orange arrow points from the text 'extract the muon magnetic anomaly' to this term. A curved arrow labeled 'by measuring' points from the text to the $\vec{\omega}_a$ term.

WHAT IS THE MAGIC MOMENTUM?

extract the muon magnetic anomaly

$$\vec{\omega}_a = -\frac{q}{m} \left(a_\mu \vec{B} - a_\mu \frac{\gamma}{\gamma + 1} (\vec{\beta} \cdot \vec{B}) \vec{\beta} + \left(a_\mu - \frac{1}{\gamma^2 - 1} \right) \frac{\vec{\beta} \times \vec{E}}{c} \right)$$

by measuring



WHAT IS THE MAGIC MOMENTUM?

extract the muon magnetic anomaly

$$\vec{\omega}_a = -\frac{q}{m} \left(a_\mu \vec{B} - a_\mu \frac{\gamma}{\gamma + 1} \underbrace{(\vec{\beta} \cdot \vec{B})}_{\sim 0} \vec{\beta} + \left(a_\mu - \frac{1}{\gamma^2 - 1} \right) \frac{\vec{\beta} \times \vec{E}}{c} \right)$$

by measuring

WHAT IS THE MAGIC MOMENTUM?

extract the muon magnetic anomaly

$$\vec{\omega}_a = -\frac{q}{m} \left(a_\mu \vec{B} - a_\mu \frac{\gamma}{\gamma+1} (\vec{\beta} \cdot \vec{B}) \vec{\beta} + \left(a_\mu - \frac{1}{\gamma^2-1} \right) \frac{\vec{\beta} \times \vec{E}}{c} \right)$$

by measuring

~ 0

~ 0

$$p = p_{\text{magic}} = \frac{mc}{\sqrt{a_\mu}} = 3.094 \text{ GeV}/c$$

WHAT IS THE MAGIC MOMENTUM?

extract the muon magnetic anomaly


pitch corrections: C_p E-field corrections: C_e

$$\vec{\omega}_a = -\frac{q}{m} \left(a_\mu \vec{B} - a_\mu \frac{\gamma}{\gamma+1} (\vec{\beta} \cdot \vec{B}) \vec{\beta} + \left(a_\mu - \frac{1}{\gamma^2-1} \right) \frac{\vec{\beta} \times \vec{E}}{c} \right)$$

by measuring



~ 0 ~ 0


$$p = p_{\text{magic}} = \frac{mc}{\sqrt{a_\mu}} = 3.094 \text{ GeV}/c$$




Injection with an offset of 77mm

“Inflector” magnet: provides field free injection path





Injection with an
offset of 77mm



Injection with an offset of 77mm

The image shows a large, circular particle accelerator tunnel, likely the Large Hadron Collider (LHC), with various components and infrastructure visible. A blue arrow points to a specific location on the tunnel's circumference. A dashed orange line traces a path around the tunnel, and a solid green line highlights a section of it. An inset image shows a close-up of a kicker magnet.



Kicker: tears muons onto orbit



The inset image shows a close-up of a kicker magnet, a device used to deflect particles. A blue beam of particles is visible, being deflected by the magnet's field. The magnet is surrounded by various components and infrastructure.


Injection with an offset of 77mm

Kicker: tears muons onto orbit




Run-1: lot of “issues” (sparks),
different settings -> datasets
Run-3a/3b: upgraded kicker cables,
allowing for larger kick

Run-2 and 3a




Run-3b, afterwards




Run-4/5/6: consistent with Run-3b

Run-1: lot of “issues” (sparks),
different settings -> datasets
Run-3a/3b: upgraded kicker cables,
allowing for larger kick


Run-2 and 3a




Run-3b, afterwards




Run-4/5/6: consistent with Run-3b





Electro Static Quadrupoles (ESQ)
vertical focusing



ESQ3


ESQ4

ESQ1

ESQ2

Electro Static Quadrupoles (ESQ)


- Run-1:** - broken HV resistors
- slower charge up
- “unstable” beam




Run-2/3, 4/5/6: fixed
(and monitored)

ESQ3

ESQ1



Electro Static Quadrupoles (ESQ)
Run-5/6: added RF




ESQ3


ESQ4

ESQ1

ESQ2



Electro Static Quadrupoles (ESQ)
Run-5/6: added RF




ESQ3


ESQ4

ESQ1

ESQ2



RF on some ESQ plates




ESQ3

ESQ4

ESQ1

ESQ2



Muon storage


cyclotron period: 149.2 ns

few 1000 muons at a time

(in 16* bunches, every ~1.2s)

Boosted muon lifetime: 64 μ s

Storage up to ~700 μ s




Muon storage

cyclotron period: 149.2 ns
few 1000 muons at a time
(in 16* bunches, every ~1.2s)
Boosted muon lifetime: 64 μ s
Storage up to ~700 μ s

e^+

μ^+





e^+

μ^+



2 straw-tracker stations


each 8 modules, 4 layers of 32 straws,
50:50 Ar:Ethane, res $\sim 100\mu\text{m}$







Time since injection: 5.0 us







Time since injection: 5.0 us




Coherent Betatron Oscillation (CBO) with ESQ RF (Run-5/6)






24 Calorimeters with 54 (9x6) Cherenkov PbF₂ crystals read out by SiPMs
- arrival time (~100ps) & energy of e⁺ (~5% at 2GeV)






24 Calorimeters with 54 (9x6) Cherenkov PbF_2 crystals read out by SiPMs
- arrival time ($\sim 100\text{ps}$) & energy of e^+ ($\sim 5\%$ at 2GeV)








Laser system for gain response calibration throughout data taking
stability 10^{-3} , rate difference 10^4



24 Calorimeters with 54 (9x6) Cherenkov PbF_2 crystals read out by SiPMs
- arrival time ($\sim 100\text{ps}$) & energy of e^+ ($\sim 5\%$ at 2GeV)









extract the muon magnetic anomaly

$$\omega_a = a_\mu \frac{eB}{mc}$$

by measuring

MEASURE: ω_a

Due to **parity violation** in muon decay, number of detected **high energy positrons** oscillates as muon **spin** points towards/away from detectors





Count positrons above an energy threshold

Counts **oscillate** at ω_a ; extract frequency from time spectrum

*for the final analysis we use an asymmetry weighted analysis

MEASURE: ω_a

Due to **parity violation** in muon decay, number of detected **high energy positrons** oscillates as muon **spin** points towards/away from detectors



Count positrons above an energy threshold



Counts **oscillate** at ω_a ; extract frequency from time spectrum

*for the final analysis we use an asymmetry weighted analysis

MEASURE: ω_a

Simplest model captures **exponential decay & g-2 oscillation**



$$N(t) = N_0 e^{-t/\tau} [1 + A \cos(\omega_a t - \phi)] \quad (\text{"5 parameter fit"})$$



MEASURE: ω_a

Simplest model captures **exponential decay & g-2 oscillation**

must account for **beam oscillations, muons losses,**
and **detector effects** ($\sim 1.6\text{ppm}$ shift in ω_a)



MEASURE: ω_a CORRECTIONS

$$\omega_a = \omega_a^m \left(1 + C_e + C_p + C_{pa} + C_{dd} + C_{ml} \right)$$

Total Run-2/3 correction was **622 ppb**, dominated by **E-field & Pitch**

MEASURE: ω_a CORRECTIONS

$$\omega_a = \omega_a^m \left(1 + \underbrace{C_e + C_p}_{\text{E-field \& Up/Down motion}} + C_{pa} + C_{dd} + C_{ml} \right)$$

E-field & Up/Down motion:
Spin precesses slower than
in basic equation

Total Run-2/3 correction was **622 ppb**, dominated by **E-field & Pitch**

MEASURE: ω_a CORRECTIONS

$$\omega_a = \omega_a^m \left(1 + \underbrace{C_e + C_p}_{\text{E-field \& Up/Down motion}} + \underbrace{C_{pa} + C_{dd} + C_{ml}}_{\text{Phase changes over each fill}} \right)$$

E-field & Up/Down motion:
Spin precesses slower than
in basic equation

Phase changes over each fill:
Phase-Acceptance, Differential
Decay, Muon Losses

Total Run-2/3 correction was **622 ppb**, dominated by **E-field & Pitch**


C_e : E-FIELD CORRECTION

The largest beam dynamics correction, Run-2/3: 378-469 ppb \pm 30-33ppb

Different methods:

- “*fast rotation*”: extract the momentum($\langle x \rangle$)-distribution from the cyclotron frequency spread
 - Fourier method
(needs correction from time-momentum correlations)
 - Binned Fit-method (CernExt)
(needs additional constraints between many bins)

- “*Tracking*”: extract the momentum($\langle x \rangle$)-distribution from betatron oscillations



C_e : E-FIELD CORRECTION

The largest beam dynamics correction, Run-2/3: 378-469 ppb \pm 30-33ppb

Different methods:

- “*fast rotation*”: extract the momentum($\langle x \rangle$)-distribution from the cyclotron frequency spread
 - Fourier method
(needs correction from time-momentum correlations)
 - Binned Fit-method (CernExt)
(needs additional constraints between many bins)

- “*Tracking*”: extract the momentum($\langle x \rangle$)-distribution from betatron oscillations

C_e : E-FIELD CORRECTION



The largest beam dynamics correction, Run-2/3: 378-469 ppb \pm 30-33ppb

Different methods:

- “fast rotation”: extract the momentum($\langle x \rangle$)-distribution from the cyclotron frequency spread
 - Fourier method
(needs correction from time-momentum correlations)
 - Binned Fit-method (CernExt)
(needs additional constraints between many bins)
- “Tracking”: extract the momentum($\langle x \rangle$)-distribution from betatron oscillations

Different “data”

- calorimeters
- Trackers
- miniSciFi (cross-checks)
new for Run-4/5/6



Minimally Intrusive Scintillating Fiber

C_p, C_{pa}, C_{ml} : PITCH, PHASE-ACCEPTANCE, AND MUON LOSS CORRECTIONS

C_p , C_{pa} , C_{ml} : PITCH, PHASE-ACCEPTANCE, AND MUON LOSS CORRECTIONS

Pitch:

Run-2/3: $170\text{ppb} \pm 10\text{ppb}$

- from the amplitude of the beam's vertical oscillation
- tracker data
- Corrected for the acceptance of the calorimeters

Run-4/5/6 closely follows the Run-2/3 analysis

C_p, C_{pa}, C_{ml} : PITCH, PHASE-ACCEPTANCE, AND MUON LOSS CORRECTIONS

Pitch:

Run-2/3: $170\text{ppb} \pm 10\text{ppb}$


- from the amplitude of the beam's vertical oscillation
- tracker data
- Corrected for the acceptance of the calorimeters

Phase-Acceptance:

Run-1:

Run-2/3: $-27\text{ppb} \pm 13\text{ppb}$

- calorimeter's phase-acceptance(x,y)



Run-4/5/6 closely follows the Run-2/3 analysis

Run-4/5/6 closely follows the Run-2/3 analysis

C_p, C_{pa}, C_{ml} : PITCH, PHASE-ACCEPTANCE, AND MUON LOSS CORRECTIONS

Pitch:

Run-2/3: $170\text{ppb} \pm 10\text{ppb}$

- from the amplitude of the beam's vertical oscillation
- tracker data
- Corrected for the acceptance of the calorimeters


Run-4/5/6 closely follows the Run-2/3 analysis

Phase-Acceptance:

Run-1:

Run-2/3: $-27\text{ppb} \pm 13\text{ppb}$

- calorimeter's phase-acceptance(x,y)



Run-4/5/6 closely follows the Run-2/3 analysis

Muon Loss:

Run-1:

Run-2/3: $0\text{ppb} \pm 3\text{ppb}$

- Muon losses reduced by an order of magnitude in Run-2 and afterwards

Run-4/5/6 closely follows the Run-2/3 analysis

C_{dd} : DIFFERENTIAL-DECAY

g-2 phase (ϕ_0) dependence due to the spread of muon lifetime in the beam

$$C_{dd} = -\frac{\Delta\omega_a}{\omega_a} = \frac{1}{\omega_a} \frac{d\phi_0}{dt} = \frac{1}{\omega_a} \frac{d\phi_0}{dp} \left(\frac{dp}{dt} \right)_{dd}$$

$(dp/dt)_{dd}$: (temporal) variation of beam averaged momentum

Run-2/3: -22 to -2ppb \pm 18ppb

Contributions:


- Beam-line effects
- momentum-orbit (p-x) correlations (from beam injection)
- longitudinal phase variations (p-t) at injection

Run-2/3:

- Some direct measurements + beam dynamics simulations

Run-4/5/6:

- same tools as Run-2/3
- A promising (tracker) data based method is under development



extract the muon magnetic anomaly

$$\omega_a = a_\mu \frac{eB}{mc}$$

by measuring


extract the **muon magnetic anomaly**



$$\omega_a = a_\mu \frac{eB}{mc}$$

by measuring



NMR: precession freq.
of protons in **B**

$$B = \gamma_p \omega_p$$








Filed Monitor: Fixed NMR Probes




Filed Monitor: Fixed NMR Probes




Trolley: Field Mapper



Filed Monitor: Fixed NMR Probes



Trolley: Field Mapper




~9000 2D field maps around
the ring

FIELD MAPS

RMS around the ring <20 ppm

take field maps every 3-5 days




The field between field maps (trolley runs) is tracked by the fixed NMR probes.


CALIBRATION

Calibrate to the Larmor frequency of shielded protons in a spherical sample: ω_p'

NMR probes are in the
“trolley’s magnetic environment”




water based calibration probe




CALIBRATION

Calibrate to the Larmor frequency of shielded protons in a spherical sample: ω'_p

NMR probes are in the
“trolley’s magnetic environment”



water based calibration probe




1) swap place in situ uncertainty: 17 ppb


CALIBRATION

Calibrate to the Larmor frequency of shielded protons in a spherical sample: ω'_p

NMR probes are in the
“trolley’s magnetic environment”



water based calibration probe




1) swap place in situ uncertainty: 17 ppb

2) correct for material effects
3) correct from cylinder -> sphere
uncertainty: 9 ppb

CALIBRATION

Calibrate to the Larmor frequency of shielded protons in a spherical sample: ω'_p


NMR probes are in the
“trolley’s magnetic environment”



Crosschecks:

- spherical water NMR probe (BNL)
- water spherical \leftrightarrow ^3He NMR
- against JPARC water NMR (CW)

water based calibration probe





in situ uncertainty: 17 ppb



- 2) correct for material effects
- 3) correct from cylinder \rightarrow sphere
uncertainty: 9 ppb

CROSS-CALIBRATIONS: EXAMPLE US/JP

Fermilab: pulsed NMR



J-PARC: continuous wave (CW)




CROSS-CALIBRATIONS: EXAMPLE US/JP



Cross-calibrated 4 times:

- 1.45T - Fermilab field
- 1.7T - MuSEUM field
- 3.0T - J-PARC field
- 1.45T - Fermilab field

Some discrepancies in the first two, good agreement (better than ~ 15 ppb) on the latest two iterations.



CALIBRATION

Calibrate to the Larmor frequency of shielded protons in a spherical sample: ω'_p

$$a_\mu = \frac{\omega_a}{\tilde{\omega}'_p} \frac{\mu'_p}{\mu_e(H)} \frac{\mu_e(H)}{\mu_e} \frac{m_\mu}{m_e} \frac{g_e}{2}$$

10.5 ppb uncertainty (hydrogen maser)
Metrologia **13**, 179 (1977)

bound state QED calc., exact

0.13 ppt uncertainty
PDG, dominated by
Phys. Rev. Lett. **130**, 071801 (2023)

22 ppb uncertainty
(Muonium hyper fine split.)
Phys. Rev. Lett. **82**, 711 (1999)

CALIBRATION

Calibrate to the Larmor frequency of shielded protons in a spherical sample: ω'_p

$$a_\mu = \frac{\omega_a}{\omega'_p} \frac{\mu'_p}{\mu_e(H)} \frac{\mu_e(H)}{\mu_e} \frac{m_\mu}{m_e} \frac{g_e}{2}$$

magnetic field seen by the muons (pointing to ω'_p)

10.5 ppb uncertainty (hydrogen maser)
 Metrologia **13**, 179 (1977) (pointing to μ'_p)

bound state QED calc., exact (pointing to $\mu_e(H)$)

0.13 ppt uncertainty
 PDG, dominated by Phys. Rev. Lett. **130**, 071801 (2023) (pointing to $\frac{m_\mu}{m_e}$)

22 ppb uncertainty (Muonium hyper fine split.)
 Phys. Rev. Lett. **82**, 711 (1999) (pointing to $\frac{g_e}{2}$)


CALIBRATION

Calibrate to the Larmor frequency of shielded protons in a spherical sample: ω'_p


$$a_\mu = \frac{\omega_a}{\omega'_p} \frac{\mu'_p}{\mu_e(H)} \frac{\mu_e(H)}{\mu_e} \frac{m_\mu}{m_e} \frac{g_e}{2}$$


magnetic field seen by the muons
 10.5 ppb uncertainty (hydrogen maser)
 Metrologia **13**, 179 (1977)
 bound state QED calc., exact
 0.13 ppt uncertainty
 PDG, dominated by
 Phys. Rev. Lett. **130**, 071801 (2023)
 22 ppb uncertainty
 (Muonium hyper fine split.)
 Phys. Rev. Lett. **82**, 711 (1999)

Run-2 and 3a



Run-3b:
upgraded kickers







ESQ4

ESQ1

ESQ3

$$\tilde{\omega}'_p = \langle \omega'_p \times M \rangle (1 + B_K + B_Q)$$

ESQ2






$$\tilde{\omega}'_p = \langle \omega'_p \times M \rangle \left(1 + B_K + B_Q \right)$$

ESQ3

ESQ1



ESQ2



$$\tilde{\omega}'_p = \langle \omega'_p \times M \rangle (1 + B_K + B_Q)$$

ESQ3

ESQ1





Run-4/5/6

- Measurements at much more positions, different kickers
- Two independent magnetometers/analysis teams
- New lab measurements for transverse model


$$\tilde{\omega}'_p = \langle \omega'_p \times M \rangle (1 + B_K + B_Q)$$

ESQ3

ESQ1



SYSTEMATIC UNCERTAINTY - WHAT TO EXPECT



Total syst. Run-1: 157 ppb

Total syst. Run-2/3: 70 ppb

TDR goal: 100 ppb


Run-1: a few “large” systematics

B_q : new measurements


C_{pa} : fixed broken hardware
improved running conditions

Run-2/3: many individual systematics
on a very similar level
(~20 to 30 ppb)

Run-4/5/6: very similar conditions
added RF system to the ESQ
-> reducing beam oscillations



Expect to publish the full dataset 2025
 ~ 2x improved precision
 likely still statistics limited



Expect to publish the full dataset 2025
 ~ 2x improved precision
 likely still statistics limited

Other Analysis:

Muon EDM:


Current best limit from BNL Muon g-2:
 $|d_\mu| < 1.8 \times 10^{-19} e \text{ cm (95 \% CL)}$

we aim to improve to $\sim 10^{-21} e \text{ cm}$
 -> *Mikio's talk on Thursday*

BSM searches:

CPT/LV & Dark Matter

THE COLLABORATION



Collaboration meeting at Argonne in Spring 2024


The Muon $g - 2$ Experiment was performed at the 326 Fermi National Accelerator Laboratory, a U.S. Department of Energy, Office of Science, HEP User Facility. Fermilab is managed by Fermi Research Alliance, 329 LLC (FRA), acting under Contract No. DE-AC02- 330 07CH11359. Additional support for the experiment was provided by the Department of Energy offices of HEP and NP (USA), the National Science Foundation (USA), the Istituto Nazionale di Fisica Nucleare (Italy), the Science and Technology Facilities Council (UK), the Royal Society (UK), the National Natural Science Foundation of China (Grant No. 11975153, 12075151), MSIP, NRF and IBS-R017-D1 (Republic of Korea), the German Research Foundation (DFG) through the Cluster of Excellence PRISMA+ (EXC 2118/1, Project ID 39083149), 340 the European Union Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreements No. 101006726, No. 734303, European Union STRONG 2020 project under grant agreement No. 824093 and the Leverhulme Trust, LIP-2021-01.



Argonne National Laboratory is a
U.S. Department of Energy laboratory
managed by UChicago Argonne, LLC.




Theory prediction is less clear now than in 2021, but we can still compare



Large discrepancy between experiment and WP (2020)

Theory prediction is less clear now than in 2021, but we can still compare




Large discrepancy between
experiment and WP (2020)

Significance for **Fermilab
alone** get to **5.0 σ**

Updated prediction considering
all available data will likely yield
a smaller and less significant
discrepancy


Theory prediction is less clear now than in 2021, but we can still compare



Include **BMW** result by swapping HVP from WP with their value

Note: BMW is currently the only full lattice calculation of HVP


Theory prediction is less clear now than in 2021, but we can still compare



Include **BMW** result by swapping HVP from WP with their value

Note: BMW is currently the only full lattice calculation of HVP

Theory prediction is less clear now than in 2021, but we can still compare



Following A. Keshavarzi at Lattice 2023...

Substitute **CMD-3** data
for HVP below 1 GeV


Cherry-picking one
experiment but gives a
bounding case

SND2k cannot be processed
in this way, but would fall
closer to WP (2020)

Disclaimer from A. Keshavarzi's Lattice 2023 talk:

- IMPORTANT: THIS PLOT IS VERY ROUGH!
- T1 White Paper result has been substituted by CMD-3 only for 0.33 \rightarrow 1.0 GeV.
 - The NLO HVP has not been updated.
 - It is purely for demonstration purposes \rightarrow should not be taken as final!

Theory prediction is less clear now than in 2021, but we can still compare



Following A. Keshavarzi at Lattice 2023...

Substitute **CMD-3** data
for HVP below 1 GeV

Cherry-picking one
experiment but gives a
bounding case

SND2k cannot be processed
in this way, but would fall
closer to WP (2020)


Disclaimer from A. Keshavarzi's Lattice 2023 talk:

IMPORTANT: THIS PLOT IS VERY ROUGH!

- T1 White Paper result has been substituted by CMD-3 only for 0.33 \rightarrow 1.0 GeV.
- The NLO HVP has not been updated.
- It is purely for demonstration purposes \rightarrow should not be taken as final!

COMPARING DATASETS: CROSSCHECKS

Datasets were taken at slightly different field settings




Example of one of the most basic “handles”:
other checks against day/night, temperature, ...

C_e : E-FIELD CORRECTION

$$\frac{\Delta\omega_a}{\omega_a} = -2 \frac{\beta_0}{cB_0} \delta E_x$$


$$C_e = -\left\langle \frac{\Delta\omega_a}{\omega_a} \right\rangle \approx 2n(1-n)\beta_0^2 \frac{\langle x_e^2 \rangle}{R_0^2}$$

Data subset 3E (Run-3)



$$\delta \approx (1-n) \frac{x_e}{R_0}$$

ω_a : STARTTIME SCANS





SYSTEMATIC UNCERTAINTY IMPROVEMENTS

Running Conditions

Syst. Measuemrents

Analysis Improvements

Temperature Control





SYSTEMATIC UNCERTAINTY IMPROVEMENTS

Running Conditions

Syst. Measurements

Analysis Improvements

Temperature Control






SYSTEMATIC UNCERTAINTY IMPROVEMENTS

Running Conditions

Syst. Measuemrents

Analysis Improvements

Temperature Control





SYSTEMATIC UNCERTAINTY IMPROVEMENTS

Running Conditions

Syst. Measurements

Analysis Improvements

Run-1 had **damaged resistors** in 2/32 ESQ leading to **unstable beam storage**
Redesign and fixed before Run-2: C_{pa} uncertainty is reduced (75 ppb \rightarrow 13 ppb)



SYSTEMATIC UNCERTAINTY IMPROVEMENTS



Running Conditions

Syst. Measurements

Analysis Improvements

Eddy currents from the kickers cause transient magnetic fields

Fiber based Faraday magnetometer





SYSTEMATIC UNCERTAINTY IMPROVEMENTS

Running Conditions

Syst. Measurements

Analysis Improvements

Mechanical vibrations of ESQ plates cause magnetic field changes.
at one location (in ESQ 4)





SYSTEMATIC UNCERTAINTY IMPROVEMENTS

Running Conditions

Syst. Measurements

Analysis Improvements

Mechanical vibrations of ESQ plates cause magnetic field changes.
at one location (in ESQ 4)





SYSTEMATIC UNCERTAINTY IMPROVEMENTS

Running Conditions

Syst. Measurements

Analysis Improvements

Mechanical vibrations of ESQ plates cause magnetic field changes.
at one location (in ESQ 4)





SYSTEMATIC UNCERTAINTY IMPROVEMENTS

Running Conditions

Syst. Measurements

Analysis Improvements

Mechanical vibrations of ESQ plates cause magnetic field changes.
at one location (in ESQ 4)




SYSTEMATIC UNCERTAINTY IMPROVEMENTS

Running Conditions


Syst. Measurements

Analysis Improvements

B_Q uncertainty: 92 ppb \rightarrow 20 ppb



vacuum sealed NMR probe



SYSTEMATIC UNCERTAINTY IMPROVEMENTS

Running Conditions


Syst. Measurements

Analysis Improvements

B_Q uncertainty: 92 ppb \rightarrow 20 ppb



vacuum sealed NMR probe




SYSTEMATIC UNCERTAINTY IMPROVEMENTS

Running Conditions


Syst. Measurements

Analysis Improvements

B_Q uncertainty: 92 ppb \rightarrow 20 ppb



vacuum sealed NMR probe



SYSTEMATIC UNCERTAINTY IMPROVEMENTS

Running Conditions


Syst. Measurements

Analysis Improvements

2 e^+ arriving at same time can be mistaken for 1:
can bias ω_a

Reduce uncertainty by:

- Improved reconstruction, correction algorithm



SYSTEMATIC UNCERTAINTY IMPROVEMENTS

Running Conditions



Syst. Measurements

Analysis Improvements

2 e^+ arriving at same time can be mistaken for 1:
can bias ω_a

Reduce uncertainty by:

- Improved reconstruction, correction algorithm



E-field correction depends on muon momentum distribution

Now include correlations between momentum & time of injection.

SYSTEMATIC UNCERTAINTY IMPROVEMENTS

